

Fermi National Accelerator Laboratory

FERMILAB-Conf-96/173-E

CDF

Inclusive Jet Cross Section Measurement at CDF

C. Pagliarone

For the CDF Collaboration

*Fermi National Accelerator Laboratory
P.O. Box 500, Batavia, Illinois 60510*

*Universita di Torino and I.N.F.N., Sezione di Trieste
Area di Ricerca, Padriciano 99
34012 Trieste, Italy*

July 1996

Published Proceedings at the *International Workshop on Deep Inelastic Scattering and Related Phenomena*, Rome, Italy, April 15-19, 1996

Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

INCLUSIVE JET CROSS SECTION MEASUREMENT AT CDF

C. PAGLIARONE

*Universita' di Torino and I.N.F.N., Sezione di Trieste,
AREA DI RICERCA, Padriciano 99,
34012 TRIESTE, Italy*

(PRESENTED ON BEHALF OF THE CDF COLLABORATION)

The CDF Collaboration has measured the inclusive jet cross section using 1992-93 collider data at $\sqrt{s} = 1.8$ TeV. The CDF measurement is in very good agreement with NLO QCD predictions for transverse energies (E_T) below 200 GeV. However it is systemically higher than NLO QCD predictions for E_T above 200 GeV.

1 Jet Inclusive Cross Section

1.1 Introduction

The measurement of the inclusive jet cross section provides a conceptually simple, but fundamental test of QCD. This measurement has, at CDF, a very good statistical precision, typically a few percent, with relatively small experimental systematic errors, about 20-30% depending on the E_T . The NLO QCD calculations have small theoretical uncertainties¹ due to the choice of renormalization/factorization scale (μ). Within the framework of conventional QCD, the study of the jet inclusive cross section is also useful to extract the strong coupling constant². Jet production at the Tevatron is dominated by gluon-gluon scattering at low E_T . Whereas at high E_T the main contribution comes from quark-quark scattering.

1.2 Data Sample and Event Selection

At CDF³ jets are reconstructed using a fixed cone algorithm which is similar to the one used in the NLO theory. The cone size used in the present measurement is $R = 0.7$ where $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2}$. CDF collected 19.3 pb^{-1} of data during the 1992-93 collider run. The measurement was performed using four different triggers with E_T thresholds of 20, 50, 70 and 100 GeV with pre-scale factors of 500, 20, 6 and 1 respectively. The data are corrected for the different trigger efficiencies. Events were required to pass the following selection criteria:

- $E_T > 15$ GeV;
- At least one reconstructed vertex with $|z| < 60$ cm;

- Missing E_T significance: $S < 6$ ($S \equiv E_T/(\sum E_T)^{1/2}$);
- Negligible energy in the calorimeter out-of-time with the $p\bar{p}$ collision.

1.3 Corrections to the Jet Cross Section

Data are corrected for detector effects using an unsmearing procedure⁴. Detector effects include energy loss due to uninstrumented regions of the detector and the smearing effects due to the finite detector resolution. The measured E_T is parametrized as a function of E_T^{True} . The E_T^{True} is defined as the sum E_T of all particles in a cone R around the jet direction. This parametrization is called *Response Function*. A very large sample of dijet events was generated and fragmented using the Feyman-Field fragmentation model, tuned to CDF data. This simulated sample was used to extract the response function. An hypothetical cross section as function of E_T is smeared by applying this response function and then fit to the data. The correspondence between smeared and unsmeared curves was used in order to derive the bin-by-bin corrections applied to the jet cross section distribution.

1.4 Results

The corrected cross-section is shown in Fig. 1.a compared to the NLO QCD predictions¹ using MRSD0' parton distribution function with a $\mu = E_T/2$. There is very good agreement between the data and NLO QCD over seven orders of magnitude. However the data deviate from the theory predictions for E_T above 200 GeV.

2 Systematic Uncertainties

The systematic errors have been evaluated by varying each of the sources of systematic uncertainty by $\pm 1\sigma$ and reapplying the unsmearing procedure, Fig 1.b. No single source of systematic uncertainty can explain the excess of events at high E_T without destroying the agreement at low E_T . Even allowing 3σ distortion does not provide a good agreement with QCD.

3 Possible Explanations of High E_T Excess

The possible explanations for this deviation include

- The choice of renormalization and factorization scale and PDFs;
- Corrections to the NLO QCD predictions, e.g. soft gluon resummation;




Figure 1: a) The inclusive Jet Cross Section with NLO QCD predictions evaluated with MRSD0' (*small figure*), comparison with other pdfs (*big figure*); b) The percentage change in the inclusive jet cross section when various sources of systematic uncertainties are changed by 1σ -standard deviation from the nominal values.

- Experimental systematics such as jet energy scale;
- New physics;

4 Conclusion

The inclusive cross section measured by CDF is in excellent agreement with NLO QCD below E_T of 200 GeV. Above 200 GeV the measured cross section begins to deviate from NLO predictions with an excess of 20-50% in the 260-360 GeV range. CDF observes a similar excess in the two-jet mass distribution and the $(\sum E_T)$ cross section. It is possible that the high E_T excess in these three measurements has a common explanation either as an experimental artifact or lack of theoretical understanding.

References

1. S. Ellis et al., *Phys. Rev. Lett.* **62**, 2188 (1989).
2. W. Giele et al., *FERMILAB Pub* **95**, 127-T (1995).
3. CDF Collaboration, F. Abe et al., *Nucl. Instrum. Methods A* **271**, 387 (1988).
4. CDF Collaboration, F. Abe et al., *Phys. Rev. Lett.* **70**, 1376 (1993).